

## Patient Instructions: Outpatient Therapy with the GliaSite® RTS

In order to release a patient undergoing radiotherapy with the GliaSite RTS, written and verbal instructions should be given to the patient and understood by the patient. These instructions are meant to provide guidance in the good practices of radiation safety and contamination prevention for these patients.

For the duration of the outpatient therapy:

1. Sleep alone.
2. Do not return to work.
3. Maintain a prudent distance from other people as much as possible (e.g. > 3 feet).
4. Keep the toilet especially clean by flushing twice after use. Men should sit during urination. Wash thoroughly after using toilet.
5. Refrain from traveling by airplane or other mass transportation.
6. Refrain from traveling by automobile except for trips to and from doctor/hospital.
7. Avoid prolonged contact with children and pregnant women.
8. Terminate breast feeding (if applicable)
9. Remember to return to hospital on scheduled Date and time: \_\_\_\_\_ AM/PM \_\_\_\_\_  
(circle one) (Date)
10. In case of emergency or if you have any questions, call \_\_\_\_\_ at \_\_\_\_\_.

I have received these instructions and fully understand them.

\_\_\_\_\_  
Patient signature                      Date

\_\_\_\_\_  
Signature of responsible  
family member or guardian                      Date

These instructions were given by:

\_\_\_\_\_  
Signature                      Date